



Fructans in *Brachypodium distachyon* as an alternative carbon sink to lignin

Nielsen, Tom Hamborg; Lundmark, Maria Therese; Ulvskov, Peter; Jørgensen, Bodil; Borkhardt, Bernhard

Publication date:
2010



Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):

Nielsen, T. H., Lundmark, M. T., Ulvskov, P., Jørgensen, B., & Borkhardt, B. (2010). *Fructans in Brachypodium distachyon as an alternative carbon sink to lignin*. Poster session presented at Synthetic Biology Workshop, Berkeley, United States.

Synthetic Biology Workshop

23-25 July 2009
Berkeley, California



INNOVATION CENTER DENMARK
SILICON VALLEY



jbei
Joint BioEnergy Institute



The Danish Council for Strategic Research
Danish Agency for Science
Technology and Innovation

Fructans in Brachypodium distachyon as an alternative carbon sink to lignin

Tom Hamborg Nielsen¹, Maria Lundmark¹, Peter Ulvskov^{1,2}, Bodil Jørgensen², Bernhard Borkhardt²

¹ Dept. Plant Biology and Biotechnology, Faculty of Life Sciences, Univ. of Copenhagen.

²Institute of Genetics and Biotechnology, Faculty of Agricultural Sciences, University of Aarhus

We present a newly started project, where the concept is to change carbon flux away from lignin and in direction of production of soluble degradable storage polysaccharides. This will be done by genetic transformation of *Brachypodium distachyon* 1) to gently reduce lignin biosynthesis by targeting DAHP synthase and 2) to introduce genes encoding fructosyl transferases, that will synthesize fructans as a carbon storage in leaves.

Along with this we will select and express genes encoding fructan hydrolases in microorganisms, in order to obtain new enzymes suitable for degrading the fructans stored in grass leaves.

Fructans in *Brachypodium distachyon* as an alternative carbon sink to lignin

Tom Hamborg Nielsen¹, Maria Lundmark¹,
Peter Ulvskov^{1,2}, Bodil Jørgensen², Bernhard Borkhardt²

(1)Molecular Plant Biology Laboratory, Dept. Plant Biology and Biotechnology, Faculty of Life Sciences, Univ. of Copenhagen
(2)Institute of Genetics and Biotechnology, Faculty of Agricultural Sciences, University of Aarhus



The concept is to ...

- change carbon flux in direction of production of soluble degradable storage polysaccharides by introducing fructosyl transferases and gently reducing lignin biosynthesis.
- express new fructanases to select enzymes suitable degrade the stored fructans

The project is a part of a new initiative "Bio4Bio".

Bio4Bio

"A strategic centre for the development and implementation of biotechnology for bioenergy" funded by Danish Council for Strategic Research

Focus: lignocellulosic plant biomass as a resource for making renewable liquid fuels.

The centre will coordinate research groups working within plant biology, biomass technology and microbiology for optimal conversion of non-food plant biomass. The research objectives are organized in the following Work Packages:

Biomass selection and development

Exploiting potential in existing material

Core technologies - including:

- Plant transformation (*Brachypodium distachyon*),
- Saccharification screening
- Characterization of plant and cell wall structural features

Protein Discovery - including:

- New Cell wall degrading enzymes and accessory proteins,
- New fructanases

Functional genomics - Including:

- Targeted approaches using plant transformation
- Production and selections of TILLING mutants

Testing and up-scaling

Fructans

Polysaccharides composed of fructose units linked by $\beta(2 \rightarrow 1)$ and $\beta(2 \rightarrow 6)$ glycosidic bonds. In many plants fructans represent a major carbon storage.

Fructans occur as different types:


- Inulin - linear fructans generally linked by $\beta(2 \rightarrow 1)$ glycosidic bonds
- Levan - linear fructans generally linked by $\beta(2 \rightarrow 6)$ glycosidic bonds
- Branched - fructans containing both $\beta(2 \rightarrow 1)$ and $\beta(2 \rightarrow 6)$ glycosidic bonds.

Examples:



Fructans are synthesized by fructosyl transferases. The transfer of a fructose from either sucrose (SST, SFT) or in between fructan chains (FFT). A range of transferases with different specificities occur and give rise to fructans with different structure and DP.


Example of activity



Fructanases

Fructans are degraded by hydrolysis by specific fructan exohydrolases (FEH)

Inulase (1-FEH) will cleave $\beta(2 \rightarrow 1)$ glycosidic bonds and whereas 6-FEH cleaves $\beta(2 \rightarrow 6)$ in levans.



Inulase is commercially available whereas 6-FEH is not

Task:

- Isolate genes encoding fructosyl transferases from *Brachypodium* and from other plant sources.

- Make constructs for overexpression of fructosyl transferases in *Brachypodium*.

- Analyze the natural fructan metabolism in *Brachypodium*.

Task:

- 1) Isolate genes encoding 6-FEH and express these in microorganisms.

Goal:


To be used for fructan degradation and as analytical tool.

Brachypodium distachyon.


Brachypodium will be used as a model plant for temperate grasses. Transformation techniques will be established to modify plants.

Temperate grasses can store large amounts of fructans in leaves, typically during cold stress and other conditions leading to sugar accumulation in the leaves. *Brachypodium* is not yet characterized in this respect.

Lignin



The entry to synthesis of lignin is catalyzed by DHAP synthase (the target enzyme of glyphosate).



Task:

- Reliable procedures for transformation of *Brachypodium* are now established

Task:

- Identify the genes encoding DHAP synthase in *Brachypodium*, and make constructs for reduced expression in *Brachypodium*.

Task & Goal :

Transformation of *Brachypodium* to obtain over-accumulation of fructans and reduced lignification in leaves.